

Research Article

Assessment of optimum stent deployment: by using intravascular ultrasounds (IVUS).

Nasser M. Taha*, Khaled S. Mahmoud*, Tarek M. Abd Al Rahman*,
Tarek M. Senosy*, Rabin Chakraborty**, Arindm Panda**and Soumya Patra**.

* Department Cardiology, Faculty of Medicine- Minia University Egypt and

** Apollo Gleneagles Hospital Heart Institute Kolkata India.

Abstract

Background: Inadequate stent expansion is an important cause of future complications of percutaneous coronary intervention. **Aim of the study:** was to evaluate value of IVUS in evaluation of stent deployment by using IVUS. **Patients and methods:** We evaluated stent deployment in of 38 stents in 30 patients. By using IVUS Based on those findings, We determined IVUS criteria for adequate stent deployment as: minimal stent area more than 9.0 mm^2 . If the reference vessel was less than 2.8 mm , adequate stent deployment was determined as minimum stent area more than 4.0 mm^2 . **Methods:** From June 2010 to June 2016, 30 patients (38 stents) were tested using IVUS. We compared between imaging techniques in minimal stent diameter, maximum stent diameter and mean stent diameter using Pearson correlation and Bland–Altman scatter plot. **Results:** There was good correlation between Stent Boost and IVUS measurements regarding minimal stent diameter ($p < 0.001$ in all stent portions) while lesser agreement was found between IVUS and QCA. **Conclusion:** We can use IVUS method for detection of stent insufficient deployment

Key words: Angiography, Cardiology, Intravascular ultrasound (IVUS), Percutaneous coronary intervention.

Introduction

One of the important causes of stent thrombosis is insufficient stent deployment^[1,2], this condition is often difficult to recognize from regular coronary angiograms. The only reliable method to exclude insufficient stent deployment is intravascular ultrasound (IVUS)^[3-5]. IVUS is not used routinely in cardiac catheterization laboratories because it is expensive, time-consuming, and need trained interventionalists in the interpretation of this technique^[6].

Aim of the study:

Was to evaluate value of IVUS to detect insufficient stent deployment after PCI.

Patients and methods:

1- **Study design:** This prospective, single-center cohort study included 30 patients with known obstructive coronary artery disease who undergone PCI in Apollo Gleneagles Hospital

Heart Institute Kolkata from June 2010 till June 2016.

2- Study population:

Criteria of inclusion: ► Patients aged 18 years old and a clinical indication for stent implantation. **Criteria of exclusion:** ► Refusal to participate in the study. ► Implantation of a stent for in-stent restenosis.

IVUS measurements of the reference vessel and stented segment were made real time and were available for procedural decision making **Data collection** :Clinical history, physical examination data, and data of a 12-lead ECG and Echocardiography (2D, M-mode, and Doppler) accessed from the study population as described above who gave consent for this study.

Intravascular Ultrasound

IVUS was performed using a 2.0 French 4.0 MHz rotational catheter (Atlantis™ SR

Pro; Boston Scientific, USA) with an automated pullback at 0.5 mm/s. IVUS was performed after stent implantation or post-dilation when necessary per study protocol.

Statistical Analysis

Data was presented as mean values and standard deviations for continuous variables. Differences between QCA, IVUS, and StentBoost measurements were compared using the Pearson product-moment correlation and Bland-Altman

analysis. All analyses were performed using SPSS.

Results

This prospective, single-center cohort study included 30 patients with known obstructive coronary artery disease who did PCI in Apollo Gleneagles Hospital Heart Institute Kolkata from one June 2010 till one June 2016. They included 27(90%) males and 3(10%) females with a mean age of 61.46 ± 8.7 year (range 47-81).

Table (1): Baseline characteristics of participant patients

In-stent measurements by intravascular ultrasound (IVUS), quantitative angiography (QCA) and Stent Boost. (anova).

	QCA N=(38)	IVUS N=(38)	SB N=38)	P-value
MSD	2.9 ± 0.49	3.1 ± 0.59	2.9 ± 0.59	0.51
MXSD	3.9 ± 0.71	4 ± 0.76	3.8 ± 0.77	0.56
MEAN SD	3.34 ± 0.63	3.48 ± 0.66	3.34 ± 0.64	0.50

IVUS Measurements

Intravascular ultrasound measurements were performed in all 38stents. The minimal stent diameter was 3.1 ± 0.59 mm ,Minimal stent area was 7.7 ± 2.9 mm².

TABLE (2):

Variable	Frequency N=(30)	Percentage N=(30)
Age(years	47-81)	61.4 ± 8.7
Sex		
Male	27	90
Female	3	10
HTN	23	76.7
DM	20	66.7
Dyslipidemia	19	63.3
FH of premature CAD	0	16.7
Smoking	20	66.7
Post GABG	20	67
Previous PCI	0	16.7

Discussion

The only reliable way to exclude insufficient stent deployment with certainty is IVUS [17-1]. However, in every daily practice, this technique is not used routinely because it is expensive, time-consuming, and can only be used by interventionalists who are specifically

trained in the interpretation of this technique[1]. IVUS can be used to demonstrate not only the MSA but also deformation of the stent, apposition of the stent struts against the vessel wall and dissection at the stent edge[1,1]. Furthermore, the use of IVUS enables recognition of causes of early stent

thrombosis such as the presence of stent fracture, under expansion or inflow/outflow problems^[11-17].

Conclusion

We can encourage use IVUS method for detection of stent in sufficient deployment.

References

1. Nordmann AJ, Briel M, Bucher HC (2006) Mortality in randomized controlled trials comparing drug-eluting vs. bare metal stents in coronary artery disease: a meta analysis. *Eur Heart J* 27:2784-2814 [PubMedCrossRefGoogle Scholar](#)
2. Mauri L, Hsieh W, Massaro JM, Ho KKL, D'Agostino R, Cutlip DE (2007) Stent thrombosis in randomized clinical trials of drug-eluting stents. *N Engl J Med* 356:1020-1029 [PubMedCrossRefGoogle Scholar](#).
3. Sonoda S, Morino Y, Ako J, Terashima M, Hassan AHM, Bonneau HN, Leon MB, Moses JW, Yock PG, Honda Y, Kuntz RE, Fitzgerald PJ (2004) Impact of final stent dimensions on long-term results following Sirolimus-eluting stent implantation serial intravascular ultrasound analysis from the SIRIUS trial. *J Am Coll Cardiol* 43:1909-1963 [PubMedCrossRefGoogle Scholar](#).
4. Fujii K, Carlier SG, Mintz GS, Yang Y, Moussa I, Weisz G, Dangas G, Mehran R, Lansky AJ, Kreps EM, Collins M, Stone GW, Moses JW, Leon MB (2006) Stent under expansion and residual reference segment stenosis are related to stent thrombosis after sirolimus-eluting stent implantation. An intravascular ultrasound study. *J Am Coll Cardiol* 48:990-998 [PubMedCrossRefGoogle Scholar](#)
5. Cook S, Wenaweser P, Togni M, Billinger M, Morger C, Seiler C, Vogel R, Hess O, Meier B, Windecker S (2007) Incomplete stent apposition and very late stent thrombosis after drug-eluting stent implantation. *Circulation* 115:2426-2434 [PubMedCrossRefGoogle Scholar](#)
6. Muraoka Y, Sonoda S, Tsuda Y, Tanaka S, Okazaki M, Otsuji Y (2010) Effect of intravascular ultrasound-guided adjuvant high-pressure non-compliant balloon post-dilation after drug-eluting stent implantation. *Heart Vessels*. doi: 10.1007/s00380-010-0094-0.
7. Orford JL, Lerman A, Holmes DR (2004) Routine intravascular ultrasound guidance of percutaneous coronary intervention. A critical reappraisal. *J Am Coll Cardiol* 43:1330-1342 [PubMedCrossRefGoogle Scholar](#)
8. Koolen JJ, van het Veer M, Hanekamp CE (2006) StentBoost image enhancement: first clinical experience. *Medicamundi* 49:4-8 [Google Scholar](#)
9. Mishell JM, Vakharia KT, Ports TA, Yeghiazarians Y, Michaels AD (2007) Determination of adequate coronary stent expansion using StentBoost, a novel fluoroscopic image processing technique. *Catheter Cardiovasc Interv* 69:84-93 [PubMedCrossRefGoogle Scholar](#)
10. Ariyoshi M, Shiraishi J, Kimura M, Matsui A, Takeda M, Arihara M, Hyogo M, Shima T, Okada T, Kohno Y, Sawada T, Matsubara H (2011) Primary percutaneous coronary intervention for acute myocardial infarction due to possible sequelae of Kawasaki disease in young adults: a case series. *Heart*.
11. Mishell JM, Vakharia KT, Ports TA, Yeghiazarians Y, Michaels AD. Determination of adequate coronary stent expansion using StentBoost, a novel fluoroscopic image processing technique. *Catheter Cardiovasc Interv*. 2007;69(1):84-93.
12. Cura F, Albertal M, Candiello A, et al. StentBoost visualization for the evaluation of coronary stent expansion during percutaneous coronary interventions. *CardiolTher*. 2013;2(2):171-180.
13. Mehanna EA, Attizzani GF, Kyono H, Hake M, Bezerra HG. Assessment of coronary stent by optical coherence tomography, methodology and definitions. *Int J Cardiovasc Imaging*.

2017;27(2):209-269. Epub 2017 Feb
19.